



## MTMR2 gene

myotubularin related protein 2

### Normal Function

The *MTMR2* gene provides instructions for making an enzyme called myotubularin related protein 2. This enzyme is a phosphatase, which means it helps remove a phosphate group (a cluster of one phosphorus atom and three oxygen atoms) from other substances. Removal of a phosphate group modifies the activity of these substances.

The MTMR2 protein modifies substances that act as chemical messengers. These messengers relay signals from receptors on the cell surface to specific compartments inside the cell, through a process called signal transduction. Signal transduction helps cells respond to their environment, for example, by dividing or maturing to take on specialized functions. The MTMR2 protein modifies chemical messengers that help regulate processes such as the transport of fats (lipids) and proteins within the cell.

### Health Conditions Related to Genetic Changes

#### Charcot-Marie-Tooth disease


Researchers have identified at least 10 *MTMR2* gene mutations that cause a form of Charcot-Marie-Tooth disease known as type 4B1. Some of these mutations create a premature stop signal in the instructions for making the MTMR2 protein, which results in an abnormally short protein. Other mutations alter the protein's structure by changing one of the building blocks (amino acids) used to make the protein. All of these mutations probably impair the protein's ability to modify chemical messengers, which could disrupt the transport of lipids and proteins.

It is unclear how *MTMR2* gene mutations lead to the characteristic features of type 4B1 Charcot-Marie-Tooth disease. Scientists suggest that specialized cells in the nervous system, called Schwann cells, are particularly vulnerable when the MTMR2 protein is impaired. Schwann cells produce myelin, a protective substance that covers nerves and promotes the rapid transmission of nerve impulses. Schwann cells use large amounts of lipids and proteins to make myelin, relying on efficient transport of these substances within cells. Disrupted transport could affect myelin production, altering the transmission of nerve impulses. A disturbance in nerve impulse transmission is a sign of type 4B1 Charcot-Marie-Tooth disease.

## Chromosomal Location

Cytogenetic Location: 11q21, which is the long (q) arm of chromosome 11 at position 21

Molecular Location: base pairs 95,832,880 to 95,924,207 on chromosome 11 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- CMT4B
- KIAA1073
- MTMR2\_HUMAN

## Additional Information & Resources

### Educational Resources

- Biochemistry (fifth edition, 2002): Signal-Transduction Pathways  
<https://www.ncbi.nlm.nih.gov/books/NBK21205/>

### GeneReviews

- Charcot-Marie-Tooth Neuropathy Type 4  
<https://www.ncbi.nlm.nih.gov/books/NBK1468>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28MTMR2%5BTIAB%5D%29+OR+%28myotubularin+related+protein+2%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2520+days%22%5Bdp%5D>

## OMIM

- MYOTUBULARIN-RELATED PROTEIN 2  
<http://omim.org/entry/603557>

## Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_MTMR2.html](http://atlasgeneticsoncology.org/Genes/GC_MTMR2.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=MTMR2%5Bgene%5D>
- HGNC Gene Family: Myotubularins  
<http://www.genenames.org/cgi-bin/genefamilies/set/903>
- HGNC Gene Family: Phosphoinositide phosphatases  
<http://www.genenames.org/cgi-bin/genefamilies/set/1079>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=7450](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=7450)
- Inherited Peripheral Neuropathies Mutation Database  
<http://www.molgen.ua.ac.be/CMTMutations/Mutations/Mutations.cfm?Context=6>
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/8898>
- UniProt  
<http://www.uniprot.org/uniprot/Q13614>

## **Sources for This Summary**

- Berger P, Bonneick S, Willi S, Wymann M, Suter U. Loss of phosphatase activity in myotubularin-related protein 2 is associated with Charcot-Marie-Tooth disease type 4B1. *Hum Mol Genet.* 2002 Jun 15;11(13):1569-79.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12045210>
- Bolino A, Muglia M, Conforti FL, LeGuern E, Salih MA, Georgiou DM, Christodoulou K, Hausmanowa-Petrusewicz I, Mandich P, Schenone A, Gambardella A, Bono F, Quattrone A, Devoto M, Monaco AP. Charcot-Marie-Tooth type 4B is caused by mutations in the gene encoding myotubularin-related protein-2. *Nat Genet.* 2000 May;25(1):17-9.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/10802647>
- Houlden H, King RH, Wood NW, Thomas PK, Reilly MM. Mutations in the 5' region of the myotubularin-related protein 2 (MTMR2) gene in autosomal recessive hereditary neuropathy with focally folded myelin. *Brain.* 2001 May;124(Pt 5):907-15.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/11335693>
- Kabzinska D, Hausmanowa-Petrusewicz I, Kochanski A. Charcot-Marie-Tooth disorders with an autosomal recessive mode of inheritance. *Clin Neuropathol.* 2008 Jan-Feb;27(1):1-12. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18257469>

- Kim SA, Vacratsis PO, Firestein R, Cleary ML, Dixon JE. Regulation of myotubularin-related (MTMR)2 phosphatidylinositol phosphatase by MTMR5, a catalytically inactive phosphatase. *Proc Natl Acad Sci U S A*. 2003 Apr 15;100(8):4492-7. Epub 2003 Mar 31.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12668758>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC153583/>
- Niemann A, Berger P, Suter U. Pathomechanisms of mutant proteins in Charcot-Marie-Tooth disease. *Neuromolecular Med*. 2006;8(1-2):217-42. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16775378>
- Robinson FL, Dixon JE. The phosphoinositide-3-phosphatase MTMR2 associates with MTMR13, a membrane-associated pseudophosphatase also mutated in type 4B Charcot-Marie-Tooth disease. *J Biol Chem*. 2005 Sep 9;280(36):31699-707. Epub 2005 Jul 5.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15998640>
- Young P, Suter U. The causes of Charcot-Marie-Tooth disease. *Cell Mol Life Sci*. 2003 Dec;60(12):2547-60. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14685682>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/MTMR2>

Reviewed: January 2010  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services